

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 036 842 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
20.09.2000 Bulletin 2000/38

(51) Int Cl.7: C12N 9/16, C12N 15/85,
A01H 5/00

(21) Application number: 99125331.1

(22) Date of filing: 20.12.1999

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• Yokota, Akiho
Ikoma-shi, Nara-pref. (JP)
• Shigeoka, Shigeru
Sakai-city, Osaka-fu (JP)

(30) Priority: 10.03.1999 JP 6289199

(71) Applicant: Nara Institute of Science and
Technology
Ikoma City, Nara Pref. (JP)

(74) Representative:
Wibbelmann, Jobst, Dr., Dipl.-Chem.
Wuesthoff & Wuesthoff,
Patent- und Rechtsanwälte,
Schweigerstrasse 2
81541 München (DE)


(54) Use of cyanobacterial fructose-1,6-bisphosphatase to improve growth of plants

(57) Using the technique of recombinant DNA, the photosynthetic ability, in other words the primary metabolism of a higher plant, was improved to enhance the growth. This improvement results in increase of harvest

potentiality and earlier harvest of crops.

A cyanobacterial fructose-1,6-bisphosphatase/se-
doheptulose-1,7-bisphosphatase is phenotypically ex-
pressed in chloroplasts of a higher plant to improve the
productivity of the higher plant.

FIG. 1



A plasmid for the incorporation into
tobacco chloroplasts

EP 1 036 842 A2

Best Available Copy

Description

BACKGROUND OF THE INVENTION

5 1. Field of the invention

[0001] This invention relates to a method for enhancing the photosynthetic activity and growth of a higher plant to increase its harvest yield and/or to enable its earlier harvest.

10 2. Description of Related Art

[0002] The development of a recombinant DNA technique has realized the incorporation of a certain exogenous gene into a higher plant and expression regulation of an existing gene therein. Only few experiments has been attempted to improve the characteristics concerning food production, such as the production yield of serials and crops. Recently, incorporation of a gene, coding enzyme participating in photosynthesis or carbohydrate metabolism, was achieved. Such gene was incorporated in anti-sense direction to inhibit the expression of the gene. The results indicated functional importance of the enzyme as a rate-determining factor of photosynthesis or carbohydrate metabolism. A Researcher in Germany have played a major role in the research.

20 SUMMARY OF THE INVENTION

[0003] Despite of it, no attempts have been performed on phenotypic expression of a certain gene in a higher plant using recombinant DNA technique to enhance the photosynthesis, which is a primary metabolism of a higher plant, and to improve its growth.

25 [0004] The object of this invention is to achieve phenotypic expression of a certain gene in a higher plant using recombinant DNA technique to enhance the photosynthesis, which is primary metabolism of a higher plant, to improve the crop productivity and yield potentiality and/or to enable the earlier harvest of the crop.

[0005] This invention provides a method for improving the productivity of a higher plant having chloroplasts by the phenotypic expression of cyanobacterial fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase in the chloroplasts.

[0006] Moreover, this invention provides a transgenic plant comprising a higher plant with a DNA fragment incorporated therein, the DNA fragment containing a base sequence coding cyanobacterial fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase.

35 [0007] The fructose-1,6-bisphosphatase (FBPase) and sedoheptulose-1,7-bisphosphatase (SBPase) in the chloroplasts of a higher plant are the key (rate determining) enzymes of a photosynthetic reductive carbon system. The activities of these enzymes are regulated by photoreduction-potentiality. On the other hand, fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase (FBPase/SBPase), derived from cyanobacterium *Synechococcus* PCC 7942 gene, is found widely in a specific type of an prokaryotic algae - a cyanobacterium. The primary structure and enzyme properties of cyanobacterial FBPase/SBPase are different from those of the FBPase or SBPase found in the chloroplasts of a higher plant. In addition, cyanobacterial FBPase/SBPase is composed of one protein, that is, a bi-functional enzyme exhibiting two kinds of enzyme activities, FBPase and SBPase.

40 [0008] FBPase-I, derived from cyanobacterium *Synechococcus* PCC 7942 gene, is a tetramer consisting of four subunits of 40kDa identical with each other. After the treatment by 1mM H_2O_2 , the purified enzyme retained more than 80% of native enzyme activity. The enzyme activity of FBPase-I was inhibited by AMP ($K_i=0.26mM$), which is a specific inhibitor of cytoplasm-type FBPase. However, it was not inhibited by fructose-2,6- P_2 . The optimum pH for the enzyme activity was 8.0 and pl value of the enzyme was 4.8. FBPase-I hydrolyzed not only fructose-1,6-bisphosphate (Fru1,6- P_2), but also sedoheptulose-1,7-bisphosphate (Sed1,7- P_2). The activities of the purified enzyme for Fru1,6- P_2 and Sed1,7- P_2 were 11.7 $\mu mol/min/mg$ protein and 12.1 $\mu mol/min/mg$ protein, respectively. The K_m alues for Fru1,6- P_2 and Sed1,7- P_2 were 52 μM and 118 μM , respectively. The enzyme activity was proved to be dependent on Mg^{2+} concentration, also equally to typical FBPase. The dose-response curve showed sigmoidal curve equally to plastid FBPase, and the $S_{0.5}$ value was shown to be $1.4 \pm 0.1 mM$. This enzyme itself was described in "Archives of Biochemistry and Biophysics, Vol. 334, No. 1, pp. 27 to 36, 1996 : Molecular characterization and resistance to hydrogen peroxide of two fructose-1,6-biphosphatase from *Synechococcus* PCC 7942".

50 [0009] The inventor incorporated fructose-1,6-bisphosphatase/ sedoheptulose-1,7-bisphosphatase, isolated from cyanobacterium *Synechococcus* PCC 7942, into a tabacco plant so that the expressed protein was transferred to its chloroplasts. The FBPase activity, the SBPase activity and the photosynthetic ability of the transgenic plant were compared to those of the wild type strain. The results measured 7 weeks after seeding showed significant increase of these activities in the transgenic plant. Furthermore, after certain period of cultivation, the plant bodies of the transgenic plant

proved to be taller than those of the wild type strain. In the transgenic plant, the areas of the blades, the diameters of the stalks, and the numbers and lengths of the roots were larger than those in the wild type strain. In addition, the contents of hexose, sucrose and starch were proved to be increased in blades, stalks and roots of the transgenic plant, compared with those of the wild type strain.

[0010] Accordingly, the photosynthetic ability of the transgenic plant, obtained by incorporation of cyanobacterial FBPase/SBPase into a tobacco plant, was improved. As the result, the ability of the transgenic plant to synthesize carbohydrate and starch is increased, and the growth was enhanced, indicating the increase of final anabolism of the transgenic plant. Therefore, incorporation of cyanobacterial FBPase/SBPase into the chloroplasts of a higher plant was proved to be a very effective technique for producing rareripe or high-yield plants.

[0011] The effect might be explained as follows. Triggered by environmental stresses, light and oxygen toxicity causes various kinds of injuries to plant bodies, resulting in a critical and limiting factor of food production. Contrary to FBPase and SBPase derived from a higher plant, cyanobacterial FBPase/SBPase is resistant against oxygen injury and thus considered to function under various environmental stresses. Moreover, a gene encoding the cyanobacterial FBPase/SBPase does not exist in higher plants, thereby eliminating the possibility of adverse effects by gene silencing.

[0012] In this invention, a vector to produce a recombinant DNA includes plasmids pBI101, pIN19 and pMSH-1. A wide variety of useful cultivated plants and woods capable of photosynthesis can be adopted as a higher plant in which the inventive recombinant DNA is incorporated. For example, the invention may be applied to serials such as maize, rice, wheat, barely, oat wheat, millet and barnyard millet, beans such as soy bean, vegetables such as potato and tomato, useful cultivated plants such as coleseed, cotton and tobacco, and trees.

[0013] An amino acid sequence may be deleted from or added to the amino acid sequence of sequence number 1, or a part of the sequence of the sequence number 1 may be substituted with another amino acid sequence in the scope of this invention, so far as the resulting peptide retains its enzymatic activity properties as fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase. Preferably, not lower than 85 percent, more preferably not lower than 95 percent of the amino acid sequence may be overlapped or identical with the amino acid sequence of the sequence number 1.

[0014] In a base sequence of sequence number 2, a base sequence referred to as base numbers from 1 to 1068 is essential for this invention, because this base sequence corresponds to a structural gene portion, that is, an amino acid sequence of the sequence number 1. In addition, a base sequence referred to as base numbers from -180 to 1170 is the most preferred embodiment of this invention.

[0015] These and other features and advantages of this invention will become apparent upon a reading of the detailed description and drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016]

Fig. 1 is a schematic view showing the structure of a plasmid incorporated into tobacco chloroplasts.

Fig. 2 is a graph showing the heights of the plant bodies of a wild type strain and a transgenic plant under water culture.

Fig. 3 is a graph showing the photosynthetic ability of a wild type strain and a transgenic plant.

Fig. 4 is a photograph showing appearance of plant bodies of a wild type strain and a transgenic plant, on the 112th day of cultivation under water culture,

Fig. 5 is a photograph showing the appearance of blades and stems of a wild type strain and a transgenic plant, on the 112th day of cultivation under water culture,

Fig. 6 is a photograph showing the appearance of roots of a wild type strain and a transgenic plant, on the 112th day of cultivation under water culture.

Fig. 7 is a graph showing contents of intermediate metabolites of a wild type strain and a transgenic plant.

DETAILED DESCRIPTION OF EMBODIMENTS

[0017] As shown in Fig. 1, tomato *rbcS* promoter, coding region of a transit peptide and cyanobacterial fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase (S.7942 FBP/SBPase) gene (*fbp-l*) were conjugated with pBI101 to construct a plasmid. The gene named *fbp-l* indicates a base sequence referred to as base numbers from -180 to 1170 of the sequence number 2, derived from cyanobacterial *Synechococcus* PCC7942. The plasmid was incorporated into *Agrobacterium tumefaciens* LBA4404, which was used for infection of leaf disk of tobacco (*Nicotiana tabacum* cv Xanthi), to incorporate *fbp-1* into a tobacco nuclear gene. After isolating the genomic DNA, incorporation of *fbp-1* was confirmed by PCR and immunoblotting methods. 7 strains of transformants (TF1-1 to TF1-7) were obtained. The chloroplasts was isolated from transformed strains (T2 generation), then expression of S.7942 FBPase/SBPase was confirmed by western blotting. Moreover, it was confirmed, by cell fractionation, that the protein expressed from

the incorporated gene is localized in the chloroplasts. (The FBPase activity, SBPase activity and photosynthetic activity)

[0018] The FBPase activity of the blades of the plant body of the transgenic tobacco cultivated 7 weeks after seeding was compared with that of the wild type strain without the incorporated gene. The result showed that the enzyme activity of the wild type strain was 1.04 ± 0.22 $\mu\text{mol}/\text{min}/\text{mg}$ chlorophyll and that of the transgenic plant was 1.82 ± 0.24 $\mu\text{mol}/\text{min}/\text{mg}$ chlorophyll. Therefore, the activity of the transgenic plant is 1.75 times higher than that of the wild type strain.

[0019] The SBPase activity was also compared. The SBPase activity of the wild type strain was 1.37 $\mu\text{mol}/\text{min}/\text{mg}$ chlorophyll and that of the transgenic plant was 2.40 $\mu\text{mol}/\text{min}/\text{mg}$ chlorophyll. Therefore, the activity of the transgenic plant is 1.75 times higher than that of the wild type strain.

[0020] The photosynthetic activity under conventional condition (360 ppm CO_2) was compared. As the result, significant difference between the wild type strain and the transgenic plant was not observed under illumination of 0, 10, 50 and 100 $\mu\text{E}/\text{s}/\text{m}^2$. However, the photosynthetic activity of the transgenic plant increased significantly, under illumination of 200 $\mu\text{E}/\text{s}/\text{m}^2$, compared with that of the wild type strain. Under 1600 $\mu\text{E}/\text{s}/\text{m}^2$ of illumination, the enzyme activity of the wild type strain was 1.24 times higher than that of the wild type strain. These results are shown in Fig. 3. (Effect of transformation upon the plant growth)

[0021] Using Hogrant medium, water culture was performed on the wild type strain and the transgenic plant. The experiment was performed under condition of 400 $\mu\text{mol}/\text{m}^2/\text{s}$, a relative humidity of 60% and a temperature of 25°C. On the 63rd day, 72nd day, 77th day, 82nd day, 85th day, 90th day, 97th day, 102nd day, 105th day, 109th day, 112th day of cultivation, the heights of the plant bodies were measured. The results are shown in Fig. 2. On the 64th day of cultivation, the height of the wild type strain was 14.0 ± 4.6 cm and that of the transgenic plant was 16.6 ± 2.9 cm. However, on the 112th day of cultivation, the height of the wild type strain was 58.3 ± 7.0 cm and that of the transgenic plant was 84.5 ± 7.8 cm, indicating significant enhancement of growth in the transgenic plants (about 1.45 times). The pictures in Fig. 4 show plant body of the wild type strain (left) and that of the transgenic plant (right).

[0022] During the whole period of growth, the blades, stems and roots of the transgenic plant grew better than those of the wild type strain. That is, the leaves are thicker with broader surface area, the stems are thicker and the number of roots are larger with each root longer. Fig. 5 is a photograph showing the appearance of the blades and stems on the 112th day of cultivation, and the wild type strain is shown in the left and the transgenic plant is shown in the right. Fig. 6 is a photograph showing the appearance of roots on the 112th day of cultivation, and the wild type strain is shown in the left and the transgenic plant is shown in the right.

(The contents of metabolic intermediates)

[0023] The contents of metabolic intermediates (hexose, sucrose, starch) were measured on upper blades (fourth blade from the top), lower blades (third blade from the bottom), stems and roots of plant bodies 12th week after seeding, for comparing the contents between the wild type strain and the transgenic plant. The results are shown in Fig. 7. The contents of metabolic intermediates in the transgenic plant increased significantly in all parts, including the upper blades, the lower blades, the stems and the roots, compared with the wild type strain. Especially, hexose and sucrose contents in the upper blades considerably increased. The accumulation of starch was observed in the lower blades. This is considered that sucrose synthesized in the upper blades was transferred into the lower blades.

[0024] As shown in these results, photosynthesis in higher plants was enhanced by this invention, increasing the production of carbohydrate and starch bio-synthesized in the transgenic plant to promote the plant growth. The dry weight of the plant bodies of the wild type strain was 14.1 ± 2.2 g at the period of flower bud production, and that of the transgenic plant was 21.0 ± 1.9 g. The dry weight of the transgenic plant increased 1.5 times compared with that of the wild type strain, indicating increase of final anabolism.

[0025] As described above, photosynthetic ability was enhanced in the transgenic plant of this invention compared with the wild type strain to improve the capability of biosynthesis of carbohydrate and starch, to promote the growth and to increase the final anabolism in the transgenic plant. Accordingly, the incorporation of FBPase/SBPase into chloroplasts of a higher plant is proved to be a very effective technique to produce a rareripe and/or high-yield crop. There has been no technique which enables production of a rareripe and/or high-yield crop using recombinant DNA technique to improve the photosynthetic ability of a higher plant, which is its primary metabolism. Therefore, this invention provides an important key technique to solve the coming crisis of food shortage.

Sequence list

<110>Applicant name: President of Nara institute of science and technology
<120>Title of invention: A method to improve productivity of higher plants
and a transgenic plant

<160>Total number of sequences: 2

<210>Sequence number: 1

<211>Sequence length: 356

<212>Sequence type: PRT

<213>Organism: Cyanobacterium Synechococcus

<220>Feature of sequence

Topology: linear

Source: fructose-1,6-bisphosphatase/sedoheptulose-1,7-

bisphosphatase derived from Cyanobacterium Synechococcus PPC

7942 gene

<400>Sequence:

1	M	E	K	T	I	G	L	E	I	I	E	V	V	E	Q	15
16	A	A	I	A	S	A	R	L	M	G	K	G	E	K	N	30
31	E	A	D	R	V	A	V	E	A	M	R	V	R	M	N	45
46	Q	V	E	M	L	G	R	I	V	I	G	E	G	E	R	60
61	D	E	A	P	M	L	Y	I	G	E	E	V	G	I	Y	75
76	R	D	A	D	K	R	A	G	V	P	A	G	K	L	V	90

	91	E	I	D	I	A	V	D	P	C	E	G	T	N	L	C	105
5	106	A	Y	G	Q	P	G	S	M	A	V	L	A	I	S	E	120
	121	K	G	G	L	F	A	A	P	D	F	Y	M	K	K	L	135
	136	A	A	P	P	A	A	K	G	K	E	T	S	I	K	S	150
10	151	A	T	E	N	L	K	I	L	S	E	C	L	D	R	A	165
	166	I	D	E	L	V	V	V	V	M	D	R	P	R	H	K	180
15	181	E	L	I	Q	E	I	R	Q	A	G	A	R	V	R	L	195
	196	I	S	D	G	D	V	S	A	A	I	S	C	G	F	A	210
	211	G	T	N	T	H	A	L	M	G	I	G	A	A	P	E	225
20	226	G	V	I	S	A	A	A	M	R	C	L	G	G	H	F	240
	241	Q	G	Q	L	I	Y	D	P	E	V	V	K	T	G	L	255
	256	I	G	E	S	R	E	S	N	I	A	R	L	Q	E	M	270
25	271	G	I	T	D	P	D	R	V	Y	D	A	N	E	L	A	285
	286	S	G	Q	E	V	L	F	A	A	C	G	I	T	P	G	300
30	301	L	L	M	E	G	V	R	F	F	K	G	G	A	R	T	315
	316	Q	S	L	V	I	S	S	Q	S	R	T	A	R	F	V	330
	331	D	T	V	H	M	F	D	D	V	K	T	V	S	L	P	345
35	346	L	I	P	D	P	K	W	R	P	E	R					356

40 <110>Applicant name: President of Nara institute of science and technology
<120>Title of invention: A method to improve productivity of higher plants
and a transgenic plant

45 <160>Total number of sequences: 2
<210>Sequence number: 2

50 <211>Sequence length: 1350
<212>Sequence type: DNA
<213>Organism: Cyanobacterium Synechococcus

55 <220>Feature of sequence

Topology: linear

Source: fructose-1,6-bisphosphatase/sedoheptulose-1,7-

bisphosphatase derived from Cyanobacterium Synechococcus PPC

7942 gene

<400>Sequence:

```
-180 CGTCGCCCCGCTCCATGCCCGCAGCTGCGCCTTTGATGCCGCGGAA -136
-135 GATATTGCCGCCAACTAACGATANNAGTCACTGCGATCGCAACTA -91
-90 AAGCCAGAGATGTGAGGAGGGGATCCGGCCTTTGGTAGACTCAAC -46
-45 TGTGGAATCCCCAGAAGCAATCATCCGTAAGGAGTCAGGACGGC -1
1 GTGGAGAAGACGATCGGTCTCGAGATTATTGAAGTTGTCGAGCAG 45
46 GCAGCGATCGCCTCGGCCCGCCTGATGGGCAAAGGCGAAAAGAAT 90
91 GAAGCCGATCGCGTCGCAGTAGAAGCGATGCGGGTGCGGATGAAC 135
136 CAAGTGGAATGCTGGGCCCGCATCGTCATCGGTGAAGGCGAGCGC 180
181 GACGAAGCACCGATGCTCTATATCGGTGAAGAAGTGGGCATCTAC 225
226 CGCGATGCAGACAAGCGGGCTGGCGTACCGGCTGGCAAGCTGGTG 270
271 GAAATCGACATCGCCGTTGACCCCTGCGAAGGCACCAACCTCTGC 325
326 GCCTACGGTCAGCCCGGCTCGATGGCAGTTTTGGCCATCTCCGAG 360
361 AAAGGCGGCCTGTTTGCAGCTCCCGACTTCTACATGAAGAACTG 405
406 GCTGCACCCCCAGCTGCCAAAGGCAAAGAGACATCAATAAAGTCC 450
451 GCGACCGAAAACCTGAAAATTCTCTCGGAATGTCTCGATCGCGCC 495
496 ATCGATGAATTGGTGGTCGTGGTCATGGATCGTCCCCGCCACAAA 540
541 GAGCTAATCCAAGAGATCCGCCAAGCGGGTGCCCGCGTCCGTCTG 585
586 ATCAGCGATGGTGACGTTTCGGCCGCGATCTCCTGCGGTTTTGCT 630
631 GGCACCAACACCCACGCCCTGATGGGCATCGGTGCAGCTCCCGAG 675
676 GGTGTGATTTCCGGCAGCAGCAATGCGTTGCCTCGGCGGGCACTTC 720
721 CAAGGCCAGCTGATCTACGACCCAGAAGTGGTCAAACCGGCCTG 765
766 ATCGGTGAAAGCCGTGAGAGCAACATCGCTCGCCTGCAAGAAATG 810
811 GGCATCACCGATCCCGATCGTGTCTACGACGCGAACGAACCTGGCT 855
```

856 TCGGGTCAAGAAGTGCTGTTTGC GGCTTGCGGTATCACCCCGGGC 900
 901 TTGCTGATGGAAGGCGTGCGCTTCTTCAAAGGCGGCGCTCGCACC 945
 946 CAGAGCTTGGTGATCTCCAGCCAGTCACGGACGGCTCGCTTCGTT 990
 991 GACACCGTTCACATGTTTCGACGATGTCAAACGGTTAGCCTGCCG 1035
 1036 TTAATTCCTGATCCCAAATGGCGGCCGGAGCGGTAGAACGGGTAT 1080
 1081 AGCTCGATCGCTTCGGTCGTTGTTTTTCAGCGAATCCATTTGCGA 1125
 1126 TCGCTTTTCAAACCCTTTTTTCGTCAACCTTCTTTAAACGGCCTC 1170

Claims

1. A method for improving the productivity of a higher plant having chloroplasts by phenotypically expressing fructose-1,6-bisphosphatase/ sedoheptulose-1,7-bisphosphatase derived from Cyanobacterium Synechococcus in said chloroplasts.
2. A method for improving the productivity of a higher plant having chloroplasts by phenotypically expressing a protein containing the following amino acid sequence (a) or (b) in said chloroplasts.
 - (a) an amino acid sequence referred to as amino acid numbers from 1 to 356 in sequence number 1 in a sequence list,
 - (b) an amino acid sequence in which a part of said amino acid sequence (a) is deleted or another amino acid sequence is added to said amino acid sequence (a) or a part of said amino acid sequence (a) is substituted with another amino acid sequence, the amino acid sequence (b) exhibiting enzyme activity as fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase.
3. A transgenic plant comprising a higher plant having chloroplasts, the transgenic plant comprising a DNA fragment incorporated in said higher plant, and the DNA fragment containing a base sequence coding fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase derived from Cyanobacterium Synechococcus.
4. The transgenic plant according to claim 3, wherein phenotypic expression of fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase is localized in said chloroplasts.
5. A transgenic plant comprising a higher plant having chloroplasts, the transgenic plant comprising a DNA fragment incorporated in said higher plant, and the DNA fragment containing a base sequence coding the following amino acid sequence (a) or (b).
 - (a) an amino acid sequence referred to as amino acid numbers from 1 to 356 in sequence number 1 in a sequence list,
 - (b) an amino acid sequence in which a part of said amino acid sequence (a) is deleted or another amino acid sequence is added to said amino acid sequence (a) or a part of said amino acid sequence (a) is substituted with another amino acid sequence, the amino acid sequence (b) exhibiting enzyme activity as fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphosphatase.
6. The transgenic plant according to claim 5 wherein said DNA fragment contains the following base sequence (c) or (d).

(c) a base sequence referred to as base numbers from 1 to 1068 in sequence number 2 in a sequence list,
 (d) a base sequence hybridizes with said base sequence (c) under stringent condition, the base sequence (d)
 encoding a protein exhibiting enzyme activity as fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphos-
 phatase.

5

7. The transgenic plant according to claim 6, wherein said DNA fragment contains the following base sequence (e)
 or (f).

10

(e) a base sequence referred to as base numbers from -180 to 1170 in sequence number 2 in a sequence list,
 (f) a base sequence hybridizes with said base sequence (e) under stringent condition, the base sequence (f)
 encoding a protein exhibiting enzyme activity as fructose-1,6-bisphosphatase/sedoheptulose-1,7-bisphos-
 phatase.

15

8. The transgenic plant according to claim 5, wherein the phenotypic expression according to said DNA fragment is
 localized in said chloroplasts.

9. The transgenic plant according to claim 6 or 7, wherein the phenotypic expression according to said DNA fragment
 is localized in said chloroplasts.

20

10. A transgenic plant of a higher plant having chloroplasts, said transgenic plant comprising fructose-1,6-bisphos-
 phatase/sedoheptulose-1,7-bisphosphatase derived from Cyanobacterium Synechococcus phenotypically ex-
 pressed in said chloroplasts.

25

11. A transgenic plant of a higher plant having chloroplasts, said transgenic plant comprising a protein phenotypically
 expressed in said chloroplasts, and the protein containing the following amino acid sequence (a) or (b).

30

(a) an amino acid sequence referred to as amino acid numbers from 1 to 356 in sequence number 1 in a
 sequence list,
 (b) an amino acid sequence in which a part of said amino acid sequence (a) is deleted or another amino acid
 sequence is added to said amino acid sequence (a) or a part of said amino acid sequence (a) is substituted
 with another amino acid sequence, the amino acid sequence (b) exhibiting enzyme activity as fructose-1,6-bi-
 sphosphatase/sedoheptulose-1,7-bisphosphatase.

35


40

45

50


55

FIG. 1



A plasmid for the incorporation into tobacco chloroplasts

FIG. 2



Comparison on the growth of plant bodies of a wild strain and transgenic plant

FIG. 3




FIG. 4




FIG. 5





FIG. 6



FIG. 7



Comparison of contents of metabolite intermediates
(hexose, sucrose and starch) produced by photosynthesis

□ Wild strain ■ Transgenic plant

THIS PAGE BLANK (USPTO)

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 036 842 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
28.08.2002 Bulletin 2002/35

(51) Int Cl.7: C12N 9/16, C12N 15/85,
A01H 5/00, C12N 15/82

(43) Date of publication A2:
20.09.2000 Bulletin 2000/38

(21) Application number: 99125331.1

(22) Date of filing: 20.12.1999

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• Yokota, Akiho
Ikoma-shi, Nara-pref. (JP)
• Shigeoka, Shigeru
Sakai-city, Osaka-fu (JP)

(30) Priority: 10.03.1999 JP 6289199

(74) Representative:
Wibbelmann, Jobst, Dr., Dipl.-Chem.
Wuesthoff & Wuesthoff,
Patent- und Rechtsanwälte,
Schweigerstrasse 2
81541 München (DE)

(71) Applicant: Nara Institute of Science and
Technology
Ikoma City, Nara Pref. (JP)


(54) Use of cyanobacterial fructose-1,6-bisphosphatase to improve growth of plants

(57) Using the technique of recombinant DNA, the photosynthetic ability, in other words the primary metabolism of a higher plant, was improved to enhance the growth. This improvement results in increase of harvest

potentiality and earlier harvest of crops.

A cyanobacterial fructose-1,6-bisphosphatase/se-
doheptulose-1,7-bisphosphatase is phenotypically ex-
pressed in chloroplasts of a higher plant to improve the
productivity of the higher plant.

FIG. 1



*A plasmid for the incorporation into
tobacco chloroplasts*

EP 1 036 842 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 12 5331

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
D,Y	TAMOI MASAHIRO ET AL: "Molecular characterization and resistance to hydrogen peroxide to two fructose-1,6-bisphosphatases from <i>Synechococcus</i> PCC 7942." ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS, vol. 334, no. 1, 1996, pages 27-36, XP002203307 ISSN: 0003-9861 * abstract; figure 5 * * page 35, left-hand column *	1-11	C12N9/16 C12N15/85 A01H5/00 C12N15/82
Y	WO 98 58069 A (MONSANTO CO) 23 December 1998 (1998-12-23) * abstract *	1-11	
A	US 5 693 507 A (DANIELL HENRY ET AL) 2 December 1997 (1997-12-02) * abstract *	1	
A	RAINES CHRISTINE A ET AL: "New insights into the structure and function of sedoheptulose-1,7-bisphosphatase; an important but neglected Calvin cycle enzyme" JOURNAL OF EXPERIMENTAL BOTANY, OXFORD UNIVERSITY PRESS, GB, vol. 50, no. 330, January 1999 (1999-01), pages 1-8, XP002148063 ISSN: 0022-0957 * page 6, left-hand column *	1	TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12N
A	DE 195 02 053 A (INST GENBIOLOGISCHE FORSCHUNG) 18 July 1996 (1996-07-18) * abstract *	1	
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 25 June 2002	Examiner Ceder, O
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 12 5331

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	<p>HARRISON E P ET AL: "REDUCED SEDOHEPTULOSE-1,7-BISPHOSPHATASE LEVELS IN TRANSGENIC TOBACCO LEAD TO DECREASED PHOTOSYNTHETIC CAPACITY AND ALTERED CARBOHYDRATE ACCUMULATION"</p> <p>PLANTA, SPRINGER VERLAG, DE, vol. 204, no. 1, 1998, pages 27-36, XP000946358</p> <p>ISSN: 0032-0935</p> <p>* abstract *</p> <p>-----</p>	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
BERLIN		25 June 2002	Ceder, O
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 12 5331

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

25-06-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9858069	A	23-12-1998	AU 735383 B2	05-07-2001
			AU 8144598 A	04-01-1999
			BR 9810171 A	06-11-2001
			CN 1267333 T	20-09-2000
			EP 1002114 A1	24-05-2000
			HU 0004228 A2	28-03-2001
			JP 2001520522 T	30-10-2001
			NO 996218 A	14-02-2000
			NZ 501781 A	27-07-2001
			PL 337581 A1	28-08-2000
			TR 9903157 T2	21-06-2000
			WO 9858069 A1	23-12-1998
			ZA 9805261 A	07-01-1999

US 5693507	A	02-12-1997	US 5932479 A	03-08-1999

DE 19502053	A	18-07-1996	DE 19502053 A1	18-07-1996
			AU 715382 B2	03-02-2000
			AU 4485896 A	31-07-1996
			CA 2209932 A1	18-07-1996
			WO 9621737 A1	18-07-1996
			EP 0802982 A1	29-10-1997
			HU 9800420 A2	29-06-1998
			JP 11503304 T	26-03-1999
			US 6245967 B1	12-06-2001

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)